

## EFLM Paper



EUROPEAN FEDERATION OF CLINICAL CHEMISTRY  
AND LABORATORY MEDICINE



Ana-Maria Simundic\*, Karin Bölenius, Janne Cadamuro, Stephen Church, Michael P. Cornes, Edmée C. van Dongen-Lases, Pinar Eker, Tanja Erdeljanovic, Kjell Grankvist, Joao Tiago Guimaraes, Roger Hoke, Mercedes Ibarz, Helene Ivanov, Svetlana Kovalevskaya, Gunn B.B. Kristensen, Gabriel Lima-Oliveira, Giuseppe Lippi, Alexander von Meyer, Mads Nybo, Barbara De la Salle, Christa Seipelt, Zorica Sumarac and Pieter Vermeersch, on behalf of the Working Group for Preanalytical Phase (WG-PRE), of the European Federation of Clinical Chemistry and Laboratory Medicine (EFLM) and Latin American Working Group for Preanalytical Phase (WG-PRE-LATAM) of the Latin America Confederation of Clinical Biochemistry (COLABIOCLI)

## Joint EFLM-COLABIOCLI Recommendation for venous blood sampling

v 1.1, June 2018

<https://doi.org/10.1515/cclm-2018-0602>

Received June 9, 2018; accepted June 10, 2018

**\*Corresponding author: Ana-Maria Simundic**, Department of Medical Laboratory Diagnostics, Clinical Hospital “Sveti Duh”, Zagreb, Croatia, E-mail: am.simundic@gmail.com, amsimundic@kbsd.hr

**Karin Bölenius**: Department of Nursing, Umeå University, Umeå, Sweden

**Janne Cadamuro**: Department of Laboratory Medicine, Paracelsus Medical University, Salzburg, Austria

**Stephen Church**: BD Life Sciences – Preanalytical Systems, Reading, UK

**Michael P. Cornes**: Department of Clinical Biochemistry, Worcester Acute Hospitals NHS Trust, Worcester, UK

**Edmée C. van Dongen-Lases**: Department of Clinical Chemistry, Academic Medical Center, Amsterdam, The Netherlands

**Pinar Eker**: Ümraniye Research and Training Hospital, Istanbul, Turkey

**Tanja Erdeljanovic**: Clinic for Otorhinolaryngology and Maxillofacial Surgery, Clinical Center of Serbia, Belgrade, Serbia

**Kjell Grankvist**: Department of Medical Biosciences, Clinical Chemistry, Umeå University, Umeå, Sweden

**Joao Tiago Guimaraes**: Department of Clinical Pathology, São João Hospital Center, Department of Biomedicine, Faculty of Medicine, Porto, Portugal; and EPI Unit, Institute of Public Health, University of Porto, Porto, Portugal

**Roger Hoke**: National Association of Phlebotomists, London, UK

**Mercedes Ibarz**: Department of Clinical Laboratory, University Hospital Arnau de Vilanova, Lleida, Spain. <http://orcid.org/0000-0003-0590-946X>

**Helene Ivanov**: Greiner Bio-One GmbH, Kremsmuenster, Austria

**Svetlana Kovalevskaya**: Clinical Laboratory Diagnostic and Pathomorphology Department, Autonomous non-profit organization of additional professional education “Institute of Laboratory Medicine”, Moscow, Russia

**Gunn B.B. Kristensen**: Norwegian quality improvement of laboratory examinations, Bergen, Norway

**Gabriel Lima-Oliveira**: Section of Clinical Biochemistry, University of Verona, Verona, Italy; and Latin American Working Group for Preanalytical Phase (WG-PRE-LATAM) of the Latin America Confederation of Clinical Biochemistry (COLABIOCLI), Verona, Italy

**Giuseppe Lippi**: Section of Clinical Chemistry, University of Verona, Verona, Italy. <http://orcid.org/0000-0001-9523-9054>

**Alexander von Meyer**: Institute of Laboratory Medicine, Kliniken Nordoberpfalz AG and Klinikum St. Marien, Weiden and Amberg, Germany

**Mads Nybo**: Clinical Biochemistry and Pharmacology, Odense University Hospital, Odense, Denmark

**Barbara De la Salle**: West Hertfordshire Hospitals NHS Trust, Operating UK NEQAS for Haematology and Transfusion, Watford, UK

**Christa Seipelt**: Sarstedt GmbH & Co.KG, Nümbrecht, Germany

**Zorica Sumarac**: Center for Medical Biochemistry, Clinical Center of Serbia, Belgrade, Serbia

**Pieter Vermeersch**: Department of Laboratory Medicine, University of Leuven, Leuven, Belgium

## Table of contents:

### Abstract

### Introduction

### Scope of the guidance

### Disclaimer

### Methodology

#### I. Pre-sampling

General considerations on appropriate mode of communication with the patient

Patient position

Step 1. Patient identification (1C)

Step 2. Verify patient is fasting and properly prepared (1B)

Step 3. Obtain supplies required for venous blood collection (2C)

Step 4. Labeling and/or identifying tubes (1C)

#### II. Sampling

Step 5. Put on gloves (1C)

Step 6. Apply tourniquet (1A)

Step 7. Select venepuncture site (1B)

Step 8. Clean sampling site (1B)

Step 9. Puncture the vein (1A)

Step 10. Drawing blood into the first tube (1A)

Step 11. Release the tourniquet (1A)

Step 12. Gently invert the tubes once immediately after collection (1B)

Step 13. Draw additional tubes following the recommended order of draw (1B)

Step 14. Remove the needle from the vein and ensure the safety mechanism is activated (1A)

Step 15. Dispose of the needle (1A)

Step 16. Bandage the puncture site (1C)

Step 17. Tell the patient to apply gentle pressure and do not bend the arm (1C)

Step 18. Invert all tubes at least 4 more times (1B)

Step 19. Remove gloves (1A)

#### III. Post sampling

Step 20. Advise the patient to rest for 5 min (1B)

#### IV. Implementation of the guidelines

Potential barriers and challenges

Framework for a successful implementation of this recommendation

### Conclusions

### References

**Abstract:** This document provides a joint recommendation for venous blood sampling of the European Federation of Clinical Chemistry and Laboratory Medicine (EFLM) Working Group for Preanalytical Phase (WG-PRE)

and Latin American Working Group for Preanalytical Phase (WG-PRE-LATAM) of the Latin America Confederation of Clinical Biochemistry (COLABIOCLI). It offers guidance on the requirements for ensuring that blood collection is a safe and patient-centered procedure and provides practical guidance on how to successfully overcome potential barriers and obstacles to its widespread implementation. The target audience for this recommendation are healthcare staff members directly involved in blood collection. This recommendation applies to the use of a closed blood collection system and does not provide guidance for the blood collection with an open needle and syringe and catheter collections. Moreover, this document neither addresses patient consent, test ordering, sample handling and transport nor collection from children and unconscious patients. The recommended procedure is based on the best available evidence. Each step was graded using a system that scores the quality of the evidence and the strength of the recommendation. The process of grading was done at several face-to-face meetings involving the same mixture of stakeholders stated previously. The main parts of this recommendation are: 1) Pre-sampling procedures, 2) Sampling procedure, 3) Post-sampling procedures and 4) Implementation. A first draft of the recommendation was circulated to EFLM members for public consultation. WG-PRE-LATAM was also invited to comment the document. A revised version has been sent for voting on to all EFLM and COLABIOCLI members and has been officially endorsed by 33/40 EFLM and 21/21 COLABIOCLI members. We encourage professionals throughout Europe and Latin America to adopt and implement this recommendation to improve the quality of blood collection practices and increase patient and workers safety.

**Keywords:** fasting; healthcare safety; patient identification; patient preparation; phlebotomy; preanalytical phase; safety needle; venous blood sampling.

## Introduction

The aim of this document is to provide a simple, condensed, risk- and evidence-based recommendation for venous blood sampling. Although several documents of the same or a similar aim and scope already exist, we believe that this document is necessary to encourage and catalyze standardization of blood collection practices across Europe and Latin America. There are several reasons behind this. A study published by EFLM WG-PRE, in 2013 showed that out of the 28 European countries

questioned, only seven had their own written nationally accepted protocols (guidelines, recommendations) for venous blood sampling [1]. Furthermore, the existing international guidelines and recommendations do not provide clear and unambiguous guidance for all steps during blood collection and some important details may not be considered. Moreover, as not all steps are equally important from the safety perspective, we believe that guidelines and recommendations should offer some level of critical assessment of potential risk associated with non-compliance. This is important to assist laboratories in prioritizing and focusing their corrective and preventive activities. Finally, the evidence behind some recommendations is not well defined or is even absent, or the quality of the evidence is not appraised or weighted.

One important aspect that has not been considered in the existing documents is how to successfully implement the recommended procedure. The current document provides a comprehensive overview of the most critical steps for a standardized blood collection procedure and practical guidance on how to successfully overcome potential barriers and obstacles to its widespread implementation.

This document is a result of the efforts of the European Federation of Clinical Chemistry and Laboratory Medicine (EFLM) Working Group for Preanalytical Phase (WG-PRE) and Latin American Working Group for Preanalytical Phase (WG-PRE-LATAM) of the Latin America Confederation of Clinical Biochemistry (COLABIOCLI) to address all the above-mentioned issues. Besides specialists in laboratory medicine, the authors of this document are representatives from national nursing associations (K.B.), hospital nurses (T.E.), phlebotomists (R.H.) and representatives from manufacturers of blood collection systems (S.C., C.S. and H.I.). Their input has been invaluable and we wish to thank them for their contribution. We encourage professionals throughout Europe and Latin America to adopt and implement this recommendation to improve the quality of blood collection practices and increase patient and worker's safety.

## Scope of the guidance

This document covers all steps of the venous blood collection procedure for in- and outpatients. The blood collection in outpatients differs from in-patients mostly in the patient preparation, patient position and physical activity prior to blood sampling. These issues are covered in the respective parts of the document. The rest of the document applies equally for in- and outpatients.

This document only applies to the use of a closed blood collection system (i.e. blood collection systems where the tube cap is not removed throughout the blood sampling process) and does not provide guidance for the blood collection with an open needle and syringe. Also, it is restricted to blood collection using needles and therefore does not cover collection from a catheter. We discourage blood sampling from an intravenous catheter, as it has been shown by many studies that catheter blood collection increases the risk of hemolysis [2–4]. In cases where catheter blood collection is the only option, care must be taken to minimize the risk of hemolysis and contamination of the sample caused by admixing of intravenous (i.v.) fluids or flushing solution (these steps are outside the scope of this document). The EFLM WG-PRE is currently working on the recommendations for catheter blood collection, to address this important issue.

Standard ISO/TS 20658:2017 “Medical laboratories – Requirements for collection, transport, receipt, and handling of samples” describes requirements that are essential for sample collection, transport, receipt and handling in an ISO 15189 setting. Our recommendation discusses best practices to fulfil those requirements, but these are neither obligatory or superior over local risk management according to recommendations in ISO 15189 and ISO 20658 [5, 6].

This document is directed to healthcare staff directly involved in blood collection (hitherto referred to in the text as a phlebotomist) as the primary target group and is limited to the venous blood collection procedure. It offers guidance on the requirements for ensuring that blood collection is a safe and patient-centered procedure. It should however be noted that all national rules and recommendations take precedence over this document if they are different in any way.

This document does not address how to obtain the consent of a patient, as this may depend on the institutional policy. Test ordering, sample handling and transport as well as collection from an unconscious patient and children are also outside the scope of this document.

## Disclaimer

Different manufacturers offer different products for venous blood collection. This document applies equally to all of them. All authors of this recommendation wish to disclose here that they do not have any preferences for the use of any particular product or any manufacturer.

## Methodology

This document has been produced by EFLM WG-PRE and endorsed by the WG-PRE-LATAM, following the identification of the critical preanalytical procedures involved in venous blood sampling [7] and is, wherever possible, consistent with Clinical and Laboratory Standards Institute (CLSI) and World Health Organization (WHO) guidelines [8, 9]. The steps in the procedure are based on the best available evidence and a consensus opinion was reached following detailed discussions and involving a mixture of stakeholders including medical and scientific laboratory specialists from 16 EFLM member countries including nurses (K.B. and T.E.), phlebotomists (R.H.), specialists in laboratory medicine and representatives of venous blood collection products manufacturers (S.C., C.S. and H.I.).

Once all the steps in the venous sampling procedure were agreed, each was graded based on a system that scores both quality of the evidence and the strength of the recommendation [10, 11]. A grading system was used as it allows a gold standard process to be established, but still leaves room for arbitrary adaptation to local requirements for the less strongly graded steps. Grading spans from 1A being the strongest and best evidenced to 2C which is very weak in both evidence and recommendation strength. The grading system is provided in Table 1. Steps and respective grades for the quality of the evidence and the strength of the recommendation are provided in Table 2. The process of grading was performed as above via discussion at a face to face meeting involving the same mixture of stakeholders stated previously. Where evidence was not available, recommendation was produced as a consensus opinion based on the expertise and experience of the group members.

A first draft of the recommendation was circulated to EFLM members for public consultation. EFLM and WG-PRE-LATAM members were invited to share this document with their members and send back their collective opinion and comments to the proposed Recommendation. Eleven out of 40 EFLM members have sent back their comments. Comments received during the public consultation and replies and rebuttal to all points raised by the national societies are available at the end of this document (Supplementary Material, Appendix 1). All comments have been taken into account during the revision of this document. A revised version has been sent for voting to all 40 EFLM and 21 COLABIOCLI members. According to the EFLM Procedure Manual, EFLM Recommendations and Guidelines have to be endorsed by more than half of EFLM Member societies, to be considered a definitive statement by the EFLM [12].

Based on the results of the voting, this document has been officially endorsed by EFLM and COLABIOCLI and is to be considered an official EFLM and COLABIOCLI statement. Voting result were as follows: 33/40 EFLM members have voted in favor of this document (Albania, Austria, Belgium, Bosnia and Herzegovina, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Ireland, Israel, Italy, Lithuania, Macedonia, Montenegro, Poland, Portugal, Romania, Russia, Serbia, Slovak Republic, Slovenia, Spain, Sweden, Switzerland, Turkey, United Kingdom and Ukraine), two EFLM members voted against (The Netherlands and Norway) and five EFLM members abstained from voting (Bulgaria, Iceland, Kosovo, Latvia, Luxemburg). All 21/21 COLABIOCLI members (Argentina, Bolivia, Brasil, Costa Rica, Colombia, Cuba, Chile, Ecuador, El Salvador, España, Guatemala, Honduras, México, Nicaragua, Panamá, Paraguay, Perú, Puerto Rico, República Dominicana, Uruguay and Venezuela) voted in favor.

The authors of this document wish to thank to all who have endorsed and supported this Recommendation.

The main parts of this recommendation are: I) Pre-sampling procedures, II) Sampling procedure, III) Post-sampling procedures and IV) Implementation.

## I. Pre-sampling

### General considerations on appropriate mode of communication with the patient

Patient communication is a key to a successful patient encounter [13, 14]. During the entire blood collection process, an empathetic and confident communication with the patient is important and should always include the following basic steps:

1. Introduce yourself, maybe also with your first name for a more personal note and explain your role within the particular health care setting.
2. After you have identified the patient correctly (see Step 1 below), explain what you will be doing, why you want to do it and what the patient has to do. Act confidently and calmly. This way the patient feels more comfortable, knowing that you are a professional and competent person.
3. Tell the patient that you have come to collect her/his blood and ask if a patient agrees to have her/his blood collected. A blood sample should never be drawn if the patient resists.
4. If asked, give a reasonable time expectation for the venous blood collection procedure itself and for the

**Table 1:** Grading recommendations used in the evaluation of available evidence.

Grade of recommendation	Clarity of risk/benefit	Quality of supporting evidence	Implications
<b>1A.</b> Strong recommendation, high quality evidence	Benefits clearly outweigh risk and burdens, or vice versa	Consistent evidence from well-performed randomized, controlled trials or overwhelming evidence of some other form. Further research is unlikely to change our confidence in the estimate of benefit and risk	Strong recommendations, can apply to most patients in most circumstances without reservation. Clinicians should follow a strong recommendation unless a clear and compelling rationale for an alternative approach is present
<b>1B.</b> Strong recommendation, moderate quality evidence	Benefits clearly outweigh risk and burdens, or vice versa	Evidence from randomized, controlled trials with important limitations (inconsistent results, methodologic flaws, indirect or imprecise), or very strong evidence of some other research design. Further research (if performed) is likely to have an impact on our confidence in the estimate of benefit and risk and may change the estimate	Strong recommendation and applies to most patients. Clinicians should follow a strong recommendation unless a clear and compelling rationale for an alternative approach is present
<b>1C.</b> Strong recommendation, low quality evidence	Benefits appear to outweigh risk and burdens, or vice versa	Evidence from observational studies, unsystematic clinical experience, or from randomized, controlled trials with serious flaws. Any estimate of effect is uncertain	Strong recommendation, and applies to most patients. Some of the evidence base supporting the recommendation is, however, of low quality
<b>2A.</b> Weak recommendation, high quality evidence	Benefits closely balanced with risks and burdens	Consistent evidence from well-performed randomized, controlled trials or overwhelming evidence of some other form. Further research is unlikely to change our confidence in the estimate of benefit and risk	Weak recommendation, best action may differ depending on circumstances or patients or societal values
<b>2B.</b> Weak recommendation, moderate quality evidence	Benefits closely balanced with risks and burdens, some uncertainty in the estimates of benefits, risks and burdens	Evidence from randomized, controlled trials with important limitations (inconsistent results, methodologic flaws, indirect or imprecise), or very strong evidence of some other research design. Further research (if performed) is likely to have an impact on our confidence in the estimate of benefit and risk and may change the estimate	Weak recommendation, alternative approaches likely to be better for some patients under some circumstances
<b>2C.</b> Weak recommendation, low quality evidence	Uncertainty in the estimates of benefits, risks, and burdens; benefits may be closely balanced with risks and burdens	Evidence from observational studies, unsystematic clinical experience, or from randomized, controlled trials with serious flaws. Any estimate of effect is uncertain	Very weak recommendation; other alternatives may be equally reasonable

(<http://www.uptodate.com/home/grading-guide#GradingRecommendations>).

**Table 2:** Venous blood sampling – the order of steps.

Step	Strength of evidence
1. Identify a patient	1C
2. Verify patient is fasting and properly prepared	1B
3. Obtain supplies required for blood collection	2C
4. Label/identify tubes	1C
5. Put on gloves	1C
6. Apply tourniquet	1A
7. Select venepuncture site	1B
8. Clean sampling site	1B
9. Puncture the vein	1A
10. Draw first tube	1A
11. Release the tourniquet	1A
12. Gently invert the tube once (one full inversion)	1B
13. Draw additional tubes following order of draw	1B
14. Remove needle from the vein and activate safety feature	1A
15. Dispose of the needle	1A
16. Bandage the puncture site	1C
17. Tell a patient to apply a gentle pressure for 5–10 min and not to bend the arm	1C
18. Invert all tubes 4 times	1B
19. Remove gloves	1A
20. Advise patient to rest for 5 min and ensure bleeding has stopped before leaving the site of venous blood collection	1B

laboratory results to be returned. Be precise in your explanations. It is increasingly common practice that only electronic order management barcodes are visible for the phlebotomist. It is therefore sometimes impossible to give a reasonable time of expectation for laboratory results if individual tests ordered are not visible for the phlebotomist. In such cases, a phlebotomist should advise a patient where to look for that information.

5. Ask the patient if they feel they have been properly informed about the procedure and if there are any further questions. Be mindful and listen to the patient's concerns. Often you will get some helpful comment on which of her/his veins are better for blood collection.
6. Ask the patient if he/she is afraid of blood collection. The evidence shows that this simple question may help identify individuals who are at increased risk of experiencing vasovagal reaction (syncope) [15]. It is also advisable to ask the patient if he/she has ever had negative experiences with phlebotomy procedures in the past, to estimate the risk of syncope, or any other risk of harm or adverse effect of blood collection. If a patient is afraid, he/she should be closely monitored during and after the blood collection, in order to prevent injuries from fall during fainting. If you

feel that the patient is nervous about the forthcoming blood collection, you can give her/him a simple task to perform, such as counting upwards or taking a deep breath before the puncture. If a patient declares to be afraid of the blood collection or if fear appears during the procedure, a patient should be instructed to lie down.

### Patient position

It has been shown that change of a body position from supine to upright and *vice versa* can dramatically affect the concentration of many laboratory parameters [16–19]. Therefore, the patient should ideally not change his/her position within 15 min prior to blood sampling. If the patient was lying down, blood sampling should be done in the lying state (this is mostly the case for hospitalized patients). Outpatients should ideally rest in a sitting position for 15 min prior to blood sampling. If a change in posture is unavoidable within this time period, it should be documented to allow correct interpretation of test results [20]. If a patient has properly rested for 15 min in the waiting area, a short walk from the waiting area to the collection area is considered to be acceptable and does not need to be documented.

### Step 1. Patient identification (1C)

- 1.1 We recommend the use of identification bracelets/bands for all inpatients.
- 1.2 All patients must be positively identified, in an active and engaging manner, by asking a patient a question: “What is your name?” and “What is your date of birth?” [21].
- 1.3 For adequate identification, at least two (patient name and date of birth) and preferably one additional identifier should be used. Additional identifiers which may be used for patient identification include:
  - address
  - health insurance number
  - patient identification number
  - ID card details or any other unique personal identifier

Understandably, the more data used to identify the patient, the smaller is the chance of patient identification errors [13].

- 1.4 The patient identity must be compared with those of the blood test request. If tubes are labeled before the

blood sampling, the phlebotomist should also make sure to compare patient identity with the tube label and ensure this way the traceability of the patient identity with the test tube label. If data obtained from the patient do not match with the data on the request form or on the tube label, blood sampling procedure must be postponed until the identification issue has been resolved.

Recommendations 1.1–1.4 are grade 1C recommendations. They must be applied to all patients and on every occasion, without exception. Although we strongly recommend that this step is executed exactly as described above, there is unfortunately a paucity of evidence for exposing a patient to harm in the case of non-compliance. However, we believe that benefits of following this procedure clearly outweigh the amount of time and effort invested to ensure compliance.

## Step 2. Verify patient is fasting and properly prepared (1B)

- 2.1 In accordance with our previously published recommendation, blood for all blood tests should be drawn in the morning (between 7 and 9 am) in a fasting state, 12 h after the last meal. Water consumption is allowed during the fasting period, but patients should refrain from alcohol for 24 h prior to blood sampling. In the morning, prior to blood sampling, patients should not drink caffeine-containing beverages (coffee, energy drinks and tea). Cigarette smoking is also not permitted in the morning before the blood sampling [22]. Chewing gum should also not be used. Morning medicine should be avoided unless it is vital for the patient.
- 2.2 We recognize that fasting requirement might pose certain logistical difficulties and find it acceptable to collect blood during the day for non-fasting patients only for emergencies or for parameters for which there is evidence that fasting is not required.
- 2.3 Patient fasting status should be verified before blood is drawn. Whenever possible, blood should not be drawn if the patient is not properly prepared (emergencies are exceptions to this rule). If blood collection is done in the non-fasting state, or a patient has not been properly prepared, this fact should be documented to allow correct interpretation of test results.
- 2.4 Intense physical activity (that exceeds normal daily activity level) should be avoided 24 h before the blood sampling.
- 2.5 Time of collection of blood for therapeutic drug monitoring (TDM) will depend on the drug and indication for testing (optimizing the drug dosage, monitoring drug adherence, adverse effects, drug intoxication, etc.). Specific recommendations for the exact time of blood sampling from the ordering physician should be followed for TDM.
- 2.6 There are other potential factors such as regular or/and recent physical activity, food intake and intake of drugs, over-the-counter medicines, food supplements and herbal preparations, etc. which are known to affect the concentration of certain analytes and it should be verified whether the patient has followed necessary instructions before blood sampling [23–25]. If some of the above issues have been identified and blood sampling cannot be postponed, the laboratory staff should wherever appropriate, document all relevant pre-analytical conditions to allow a correct interpretation of test results.
- 2.7 Additional collections during the day may be advisable for tests with circadian variations. Specific recommendations from the ordering physician for the exact time of blood sampling for these tests should be followed.

The postprandial response to food and drink depends on various non-modifiable (age, gender, genetic background, blood group, etc.) and modifiable factors. Modifiable factors are diet [26–29], intake of drugs, over-the-counter medicines, food supplements and herbal preparations [30], lifestyle, physical activity, such as diving, marathon, strenuous exercise, and some other activities [31–33], body weight, smoking, alcohol consumption, etc. To limit the variation in postprandial response as a consequence of inter-individual heterogeneity the EFLM WG-PRE has in 2014 published a recommendation on how to standardize the definition of fasting requirements [22]. The above requirements are fully in line with this recommendation.

Physical activity is one very important modifiable factor which is known to exert both acute and chronic effects on human metabolism and blood composition. Whereas chronic effects of sport may be considered as adaptation of human organism, the acute effects may be obviated by avoiding intense physical activity 24 h prior to blood collection.

## Step 3. Obtain supplies required for venous blood collection (2C)

This section focuses mostly on blood sampling in an outpatient clinic and not so much in a hospitalized ward with bedridden patients.

- 3.1 Venous blood collection should be performed in a clean, quiet and private environment. The blood collection area may contain pictures with relaxing landscapes on the walls, to make the space more comfortable.
- 3.2 Dedicated venous blood collection chairs and/or bed should be in place as well as a chair for the phlebotomist. The armrests of the chair should be adjustable to enable the optimum position for blood collection to be obtained. If a dedicated venous blood collection chair is not available the chair must have arm rests to prevent patients from falling if they feel faint [8, 9, 34].
- 3.3 Hand sanitizing or washing areas with soap and/or appropriate sanitizers and paper towels should be available and accessible to ensure proper hand hygiene.
- 3.4 Patient sample collection facilities should be separated from reception/waiting areas to ensure patient privacy. Patient privacy should be ensured throughout the entire blood sampling procedure. We do recognize that conditions may differ in outpatient and inpatient settings and for inpatients with different clinical conditions. However, care should be taken to ensure that blood sampling is always done with respect to patient privacy.
- 3.5 Equipment and supplies should be available in sufficient quantities and appropriate for their intended use in the venous blood collection process. Available equipment may include:
  - utility cart
  - blood collection trays
  - gloves
  - blood collection system with safety features (needles and holders, or needles with integrated holders)
  - blood collection tubes (a full range of tubes with different volume, within the expiry date)
  - tourniquet (preferably single use)
  - antiseptics to clean the puncture site
  - bandages
  - gauze pads
  - sharps bin
  - sample mixer
  - leak proof transportation bags
- 3.6 All required materials must be assembled prior to venous blood collection and per requested tests. The workplace should be arranged so that a phlebotomist can reach all necessary supplies without leaving their place.
- 3.7 Equipment should be properly maintained and kept clean.
- 3.8 A stock management system should be in place to ensure supplies are used before expiration.
- 3.9 Needle, holder and the blood tube make together an integral blood collection system. Only individual components of the same manufacturer should be used as a part of the blood collection system. Whereas manufacturers ensure the full compatibility between the components of their system, individual components from different manufacturers should never be used together, as their combinations are not validated for the intended use and may compromise patient and healthcare worker safety [35]. If for whatever reasons, this requirement cannot be fully respected and individual components from different manufacturers need to be used together (e.g. special blood drawing tubes are not available by the main company whose tubes are in use in the particular institution), serial venepunctures to safeguard single manufacturer compatibility of blood collection system components are not justified.
 

Storing tubes under conditions not consistent with the manufacturer's recommendations can affect the draw volume, as well as the stability of gels and additives. Environmental factors such as temperature, humidity, altitude and light exposure can have a significant impact on the quality of the blood collection equipment. Pre-evacuated blood collection tubes which are beyond the expiry date have a decreased vacuum which may lead to drawing a less than optimal volume of blood and lead to an improper blood to additive ratio [36, 37]. Moreover, expired tubes may suffer from some chemical deterioration of the tube additive. To ensure sample quality, blood collection tubes should be discarded after their expiration date.

Recommendations listed under 3.1–3.8 are grade 2C recommendations (weak recommendation, low quality evidence). We were unable to find any firm evidence besides manufacturer's recommendations, one study in humans and one veterinary study [36, 37] to support the above listed recommendation.

#### Step 4. Labeling and/or identifying tubes (1C)

- 4.1 Tube labeling or tube identification (for pre-labeled tubes) must be done in the presence of the patient. Otherwise, there is a risk that the tube will be left unlabeled and possibly incorrectly identified. The choice about whether to label or identify tubes before or after blood collection should be based on a prospective

risk analysis of the venous blood collection process in each institution.

- 4.2 Each institution should have a standard written procedure to which all personnel should adhere.
- 4.3 Essential information about the sample and the patient must be registered within the laboratory in such a manner that the tube is traceable and unambiguously linked to the patient, collected sample, test request, requestor and phlebotomist. These data include but are not limited to:
  - identification of a requestor, i.e. authorized (under national law) person to order blood test
  - patient's full name
  - patient's date of birth
  - patient's address (home address or hospital department for inpatients)
  - unique sample identification number
  - date and time of sampling
  - identification of phlebotomist
- 4.4 A minimum of two independent identifiers (patient's full name and date of birth) and preferably three (two above plus additional one), e.g. unique sample identification number, should be used to identify the tube. It is not essential that all the above listed data are recorded on the blood tube. If not on the tube, this information must be documented in paper records or linked to the laboratory information system and easily retrievable.

## II. Sampling

### Step 5. Put on gloves (1C)

- 5.1 A new pair of gloves should always be worn to protect the patient and the staff performing the venous blood sampling.
- 5.2 Hands should be cleaned to minimize the risk of transmitting the infection during glove removal, but also to reassure the patient, before putting on gloves.

Unfortunately, although we consider this a strong recommendation, we were unable to find high quality evidence to support it. A recent Cochrane Database Systematic Review has shown that the role and level of protection of personal protective equipment is still unclear [38]. Nevertheless, given the potential associated risk, until proven otherwise, we recommend that gloves are used to protect the patient and the healthcare worker. In the event of a needle-stick injury, gloves act as a barrier or protection to minimize the amount of blood that might be transmitted during

the needlestick injury [39, 40]. Given the fact that a substantial proportion of healthcare staff directly involved with blood collection has at some time point been exposed to a needle-stick injury during their working time, wearing gloves sounds like a reasonable infection prevention measure [41, 42]. The evidence also shows that the use of sterile gloves during blood collection for blood culture reduces the risk of contamination of the sample [43, 44]. In addition, apart from being exposed during the needlestick injuries, venous blood sampling is always associated with a risk for blood contact and contamination during the procedure. There is evidence showing that this risk is reduced using gloves [45, 46]. It has been shown that hand cleansing is the key to the reducing the risk of the infection of the healthcare staff and cross-transmission of antimicrobial resistant pathogens. Moreover, proper hand cleaning and wearing gloves protects the patient against infections [47]. Unfortunately, the evidence shows that gloves are not widely used among healthcare workers [48].

CLSI GP41-A7 guidelines recommend putting gloves on after applying a tourniquet. However, there is evidence that the time of tourniquet application may then be longer than 1 min if this CLSI recommended procedure is followed [49]. Therefore, to reduce prolonged blood stasis we suggest the gloves are put on prior to tourniquet application.

- 5.3 Assemble the needle a) and the holder (if not already pre-assembled) or b) with an integrated holder with the blood collection tube (for users of the blood collection systems with aspiration technique).

### Step 6. Apply tourniquet (1A)

The tourniquet is conventionally defined as a constricting or compressing (elastic) device, which can be used to limit venous circulation to an extremity (usually an upper arm) for a limited period of time. In the absence of some other device which may be used to make veins visible, the use of the tourniquet may be helpful, especially in those patients with small or scarcely visible veins.

- 6.1 However, we recommend that blood collection is done preferably without tourniquets (especially in patients with prominent veins) and that tourniquets are used only when necessary. In the case when tourniquet is used, a phlebotomist should make sure that the total tourniquet time is up to 1 min.
- 6.2 The tourniquet should be applied approximately one hand width (7.5 cm) above the anticipated puncture

site and should be tight enough to stop venous but not arterial blood flow.

- 6.3 We recommend that disposable tourniquets are used to minimize the risk of infection and cross-contamination of patient and healthcare staff.

Evidence shows that reusable tourniquets can be colonized with multiresistant microorganisms and may thus serve as a reservoir and source of transmission of various pathogens to hospitalized patients [50–52]. Reusable tourniquets may even be contaminated with methicillin-resistant *Staphylococcus aureus* (MRSA) and thus pose a great risk for patients and healthcare staff. Given the risk associated with the use of reusable tourniquets and the quality of available evidence, we have graded this recommendation as 1A. Unfortunately, disposable tourniquets are not widely used, especially in some developing or non-developed countries [53]. Hospital management should be made aware of the risk associated with the use of reusable tourniquets and potential benefit of the use of disposable tourniquets for the safety of the patients and healthcare staff.

- 6.4 To minimize the risk of venous stasis, especially if multiple tubes are to be drawn, instead of tourniquets, vein illumination devices may be used to locate the veins. This is especially useful in patients with difficult veins. It has been shown that vein illumination devices may serve as an useful alternative for tourniquets to avoid venous stasis and subsequent alterations of the concentration of various biochemistry, hematology and coagulation parameters in the blood [54–56]. The use of vein illumination devices may be a valuable perspective for the future, although more clinical evidence is necessary before widespread implementation can be recommended.


- 6.5 Warn the patient not to clench or pump the fist. Fist clenching and pumping may cause pseudohyperkalemia and alterations of some other biochemistry and hematology parameters [57–62].

### Step 7. Select venepuncture site (1B)


- 7.1 To select the venepuncture site, the patient's arm should be stretched in a downward position.
- 7.2 If available, the most prominent veins in the cubital fossa (i.e. cephalic, basilic, median cubital and median antebrachial veins) should be the first choice (Figure 1). The cubital vein is the most preferable choice, as it is usually the most prominent, does not roll under the skin and can be found in the same place in most patients.
- 7.3 Only if main veins are unavailable then dorsal hand veins may be used as an alternative.
- 7.4 Blood collection from the veins in the wrist is discouraged.
- 7.5 Palpation of the vein could help in the assessment of the appropriate venepuncture site.

Cross-sectional graphic presentation of the cubital fossa is depicted in Figure 2. Understanding the anatomy of this region helps reducing the risk of injuries during the blood collection procedure.

- 7.6 Do not collect blood from previously placed peripheral venous catheters, hardened veins, artero-venous shunt, from the sites of hematoma, inflammation or swelling, from an arm with a vascular graft, paretic arms or arms with lymphatic drain disorders.
- 7.7 Make sure to document when alternate venepuncture sites (e.g. veins in hand and foot, or any other than the above-mentioned sites) are used.



**Figure 1:** The most frequent variations of the veins of the forearm. Reprinted from [63] with kind permission of the Elsevier GmbH.



**Figure 2:** Topographic anatomy of the cubital fossa (cross-section at the elbow).

Vessels: CV, cephalic vein; RA, radial artery; BV, basilic vein; tendons:  $\alpha$ , biceps brachii tendon;  $\beta$ , triceps brachii tendon; nerves: a, lateral antebrachial cutaneous nerve; b, medial antebrachial cutaneous nerve; c, median nerve; d, ulnar nerve; e, posterior lateral antebrachial nerve; f, radial nerve; muscles and bones: 1, brachioradialis; 2, brachialis; 3, pronator teres; 4, trochlea (humerus); 5, olecranon (ulna); 6, anconeus. Reprinted from [59] with kind permission of the Croatian Society of Medical Biochemistry and Laboratory Medicine.

Recommendations 7.1–7.7 are grade 1B recommendations. They must be applied to all patients and on every occasion, with no exception.

Selecting the best vein and recognizing the most appropriate site to insert the needle for venous blood collection is important for sample quality, patient satisfaction, to avoid nerve damage, to avoid arterial puncture, for the ease and speed of collection and ultimately for a successful blood collection procedure [59]. There is ample evidence demonstrating that blood collection procedures may cause some serious injuries in the case of failure to find an appropriate vein for performing the venous blood collection [64, 65].

### Step 8. Clean sampling site (1B)

8.1 The selected venepuncture site should be cleaned with 70% ethyl alcohol or any other appropriate disinfectant prior to blood sampling to prevent contamination with skin pathogens. Cleaning should be performed with one wipe and the selected site should be left to dry. Do not wipe the sampling site with the same gauze twice.

8.2 For blood culture collection, we recommend to adhere to the instructions provided by the Hospital Department of Microbiology and/or to the information provided by the disinfectant manufacturer. Cleaning the sampling site by disinfecting twice using separate gauze pads seems advisable. Let the disinfectant dry for at least 60 s [66, 67].


8.3 Do not touch the disinfected site after the cleaning.

Contamination of blood, by the normal flora of skin, during the blood collection procedure has been demonstrated to occur if the venepuncture site has not been properly cleaned [68, 69]. Cleaning is therefore of utmost importance if blood is collected for blood culture.

Alcohol evaporates quickly and already within 10 s the amount of alcohol is reduced by half of the initial amount [70]. Although the failure to let the alcohol dry may indeed cause an itchy sensation in some patients, it will not compromise the blood collection procedure and the quality of the sample. It has been shown that the presence of alcohol (in case the venepuncture site was not let to dry) on the collection site is not a source of spurious hemolysis [71]. Moreover, under ideal blood collection conditions, the use of ethanol before venous blood collection does not interfere with blood alcohol measurement [72]. Nevertheless, to avoid a risk of false-positive alcohol results, we suggest that in collections of blood samples for forensic alcohol testing the alcohol should be left to dry before performing a venous blood collection. Alternatively, non-alcoholic antiseptic cleanser approved for use by the institution may be used to avoid the risk of contamination.

### Step 9. Puncture the vein (Figure 3) (1A)

- 9.1 Puncture the vein with the bevel up, as it minimizes the pain and reduces the risk for perforation of the back wall on the vein.
- 9.2 Prevent the rolling veins by extending the patient's skin.
- 9.3 Insert the needle longitudinally into the vessel with determination and prudence at an approximately 5–30 degree angle depending on the vein's depth so that at least 0.5 cm of the needle is inserted into the vessel.
- 9.4 Hold the tube holder steady and by supporting your hand against the patient's arm. Ensure that the patient's fist is open and not clenched when blood comes [8, 9, 73].



**Figure 3:** Needle should be inserted into the vessel at an approximately 5–30 degree angle, depending on the vein's depth. (A) Inserting the needle for the users of pre-evacuated tubes and (B) inserting the needle for the users of blood collection systems using the aspiration technique.

- 9.5 If a vein cannot be located, a slight repositioning of the needle (by moving the needle backward or forward) may help to find the vein.
- 9.6 The use of sharps device with flash visualization may be helpful, especially with non-experienced staff, or in children and patients with difficult veins. These devices provide a visible venous flash when the needle is connected to the vein (Figure 4).



**Figure 4:** Blood collection device with flash visualization (butterfly – left, needle with a visible venous flash space – right).

### Step 10. Drawing blood into the first tube (1A)

10.1 Draw the blood by a) inserting the tube in the holder so that the cap is perforated and the blood is drawn (vacuum technique) or b) withdrawing the plunger slowly (aspiration technique). Follow the EFLM recommended order of draw [74]. As blood collection techniques may differ with respect to the manufacturer, specific recommendations of the manufacturer should always be followed, along with the recommendations in this document, during blood collection.

The recommended order of draw is as follows:

1. Blood culture tube
2. Citrate tube
3. Plain tube or tube with clot activator
4. Heparin tube
5. EDTA tube
6. Glycolysis inhibitor tube
7. Other tubes

10.2 When coagulation tube is collected as the first or the only tube

- and a straight needle is used for blood collection, no discard tube is needed [75, 76]
- and a winged blood collection set (butterfly devices) is used, a discard tube must be collected to prevent underfilling of the tube with subsequent bias in test results [8]

10.3 Ensure that tubes are fully filled (e.g. up to the indicated level on the tube). Underfilling of the tubes (tubes filled with less than 90% of draw volume) is strongly discouraged and should be avoided.

Although some would argue that incorrect order of draw when using closed blood collection systems is not the source of contamination [77, 78], there is firm evidence showing that contamination still occurs more commonly than might be expected and can be difficult to identify [79–82]. This is probably because venepuncture is not always performed under ideal conditions. There are still clinical settings such as emergency departments, where blood sampling is performed in less than ideal conditions and where only a minor proportion of blood collections is performed using the conventional manufacturer prescribed closed collection technique [83]. Given the reasons explained above, and because there is no obvious disadvantage in following the order of draw, we recommend that the order of draw is followed without exceptions during every blood collection.


### Step 11. Release the tourniquet (1A)

- 11.1 The tourniquet should be removed as soon as the blood flows into the first tube.
- 11.2 If the blood collection is unsuccessful, the tourniquet should be released and blood collection should be done on an alternative site.

Tourniquets cause a temporary occlusion of veins and temporary venous stasis. If applied for a long period of time (longer than 1 min) a tourniquet induces a substantial variation of blood composition, due to extravasation of water and small molecules such as ions from the vessel into the subendothelial space. During that process, large molecules such as lipoprotein particles, proteins and protein-bound substances, cells and coagulation factors remain within the vessel, so that their concentration progressively increases. Most of these changes are negligible within 1 min of the application of the tourniquet, but can become clinically significant afterwards [84–86].

### Step 12. Gently invert the tubes once immediately after collection (1B)


- 12.1 Mix all tubes once immediately after the blood has been drawn. Any delay may affect the quality of the sample.
- 12.2 Mix each tube gently by inverting it once, before collecting the next tube. One inversion involves turning the tube vertically for 180° and putting it back to the starting position (Figure 5).



**Figure 5:** One mixing cycle. One inversion involves turning the tube vertically for 180° and putting it back to the starting position. Reprinted from [25] with kind permission of the Croatian Society of Medical Biochemistry and Laboratory Medicine.

- 12.3 The dominant hand should be used to hold the needle and a holder in place throughout the collection to maintain control. Also, the hand should not be changed during the drawing of the additional tubes (Figure 6).
- 12.4 Avoid vigorous mixing of the specimens (e.g. shaking) to prevent blood cell injury, hemolysis, platelet activation or blood clotting [87].
- 12.5 The use of automated mixing tables/devices is recommended as it enables immediate mixing of samples without engaging a phlebotomist.

Appropriate mixing of the blood tube after the blood has been drawn is an important step which ensures that tube additive (anticoagulant, clot activator, etc.) is adequately mixed, blood samples are homogenous, and sample quality and integrity are maintained. We are aware that manufacturers are providing their specific recommendations on the number of inversions for a particular tube, i.e. that



**Figure 6:** Gently invert the tube once immediately after collection. Hold the needle with a dominant hand. Do not change hands during mixing and drawing of the additional tubes. (A) mixing the tube for the user of pre-evacuated tubes and (B) mixing the tube for the users of blood collection systems using the aspiration technique.

tubes should be gently inverted at least 5 to 10 times, depending on the tube type [8, 88, 89].

Over the past few years there has been a debate about whether mixing does or does not affect the quality of the sample. Some studies have shown that failure to mix the primary blood tube most probably will not introduce a bias in many test results. The explanation for these observations could be that blood turbulence that is caused by standard vacuum pressure inside the primary tubes is sufficient, in itself, to provide both solubilization, mixing and stabilization of additives and blood during venepuncture [90–92]. It could certainly be that under optimal conditions mixing the tube after venous blood collection might not be mandatory [93–95]. However, in some borderline conditions and circumstances, the failure to mix the tube may affect the quality of the sample and, for example, lead to sample hemolysis or clotting. Given the reasons explained above we strongly recommend that tube mixing is done always without exception.

In cases when more than one tube needs to be collected, mixing the first tube and putting the next tube into the holder at the same time is practically impossible, if a phlebotomist holds a holder with one hand and is mixing the tube with another hand. If a phlebotomist chooses to first mix one tube (for example, for 10 times) and only after that to leave this tube, take the next one and insert it into the holder, the average time necessary to complete the mixing and put on the next tube would be at least 15 s (unpublished observations). If multiple tubes need to be drawn, the total time during which a patient has a needle in his/her vein might be substantially prolonged. To overcome this and ease the patient discomfort, while not significantly compromising the quality of the samples, we herein recommend that if multiple tubes are to be drawn, each tube is mixed by only one full inversion and only when all tubes are collected and needle is removed from the patient vein, all tubes are mixed for additional 4 times (see Step 18).

### **Step 13. Draw additional tubes following the recommended order of draw (1B)**

- 13.1 Draw all subsequent tubes and gently mix each tube once (one full inversion), as explained in the previous step (see Step 12).
- 13.2 Draw tubes in the recommended order of draw (See Step 10).

### **Step 14. Remove the needle from the vein and ensure the safety mechanism is activated (1A)**

After disconnecting the last tube put a gauze pad on the venous blood collection area, without applying the pressure. Gently remove the needle trying not to cause any injury and press the puncture site with the gauze pad to avoid bleeding.

There are safety engineered blood collection devices on the market that may differ in the way they are activated (e.g. while the needle is still inside the vein, or after the needle has been removed from the vein). In accordance with the European Directive 2010/32 EU we recommend that only safety engineered blood collection devices are used to prevent exposure of healthcare workers and patients to a contaminated needle [96]. Manufacturers' recommendations should be followed depending on the device used.

### **Step 15. Dispose of the needle (1A)**

- 15.1 Immediately after the safety mechanism has been activated, the used blood collection device should be disposed into a puncture-resistant sharps container.
- 15.2 Sharps containers should be within arms length. Walking to sharps container is not an acceptable practice.

### **Step 16. Bandage the puncture site (1C)**

- 16.1 Check that the bleeding has stopped. Treat the wound by applying a patch or a bandage by placing an adhesive tape tight over a dry pad/gauze square.

### **Step 17. Tell the patient to apply gentle pressure on the puncture site and not to bend the arm (1C)**

- 17.1 Patient should be advised to apply gentle pressure on the puncture site and not to bend the arm, in order to minimize the risk of hematoma or prolonged bleeding.
- 17.2 Elevating the arm may be useful to stop bleeding from the puncture site.

A gentle pressure on the puncture site should be applied until the bleeding has stopped, which is usually a period of up to 2 min for routine draws and up to 10 min for patients on anticoagulation. If the cubital vein was punctured, the patient's arm should

be straight. Although one study in Denmark found no difference in the risk of bruising irrespective to whether the arm was bent or not [97], many studies have shown that bending the arm can cause a hematoma [98, 99]. Also, it has been demonstrated that a failure to apply pressure until the bleeding has stopped may increase the incidence and severity of bruising [100].

### Step 18. Invert all tubes at least four more times (1B)

- 18.1 After removing the needle from vein and activating the safety mechanism in place, invert all tubes at least 4 more times, so that a total number of inversions is five, i.e. once immediately after the tube has been filled and remaining 4 times, once all tubes have been collected (after removing the needle from vein). Ideally, the number of full rotations should correspond to manufacturers' instruction. For information about the proper mixing procedure please refer to Step 12.
- 18.2 If only one tube is collected invert it 5 times directly after collection.
- 18.3 After the mixing procedure all the tubes should be left in the upright position prior to further processing.

### Step 19. Remove gloves (1A)

- 19.1 As used gloves might be contaminated with body fluids and/or microorganisms, we recommend that gloves are changed after every venous blood collection.
- 19.2 We recommend that following procedure is used for glove removal: remove one glove and turn it inside out (Figure 7, left), enclosing the first glove by rolling the second glove over it (Figure 7, right).
- 19.3 Discard the gloves and clean your hands [101].

## III. Post sampling

### Step 20. Advise the patient to rest for 5 min (1B)

- 20.1 Advise the patient to rest for 5 min or wait until the bleeding has stopped (if longer than 5 min) before leaving the blood collection area.
- 20.2 Be empathetic and ask a patient how he/she feels before leaving the blood collection facility. This may help identify patients who are at risk of experiencing dizziness or even syncope.
- 20.3 Thank the patient and leave her/him with the assurance that she/he will obtain his laboratory results as soon as possible. If asked about the exact time for the laboratory results to be returned, either inform a patient about it or advise a patient where to look for that information (see Pre-sampling, under point 4).

With this step, we want to draw attention to the period after the blood sampling, during which patients may feel dizzy, or even faint, due to a vasovagal syncope. There are patients who are afraid of needles or feel discomfort when seeing blood. Such patients, especially young ones may in some circumstances even experience syncope during or immediately after the blood collection [102, 103]. Syncope during or after the blood collection may occur as a result of either anxiety, or a sudden relief from anxiety, when a patient no longer feels threatened [104]. Therefore, to make sure that patient is well and that no acute complications have occurred, we suggest that a patient is advised to rest for at least 5 min or longer until the bleeding has stopped, in the blood collection area or in the waiting room. Preferably, the patient should be monitored by authorized personnel, or left to rest unsupervised and advised to inform the staff or ask for help if in need for any assistance. Although we recognize that the majority of patients do not suffer



**Figure 7:** Removing the gloves: remove one glove and turn it inside out (left), enclosing the first glove by rolling the second glove over it (right).

from anxiety or dizziness post phlebotomy, we also believe that a benefit of complying to this step has an obvious benefit which outweighs possible difficulties in meeting this recommendation.

As already explained earlier (under heading: Patient Communication), empathetic and confident communication with a patient is very important. Assessing the degree of fear of blood collection may help identify patients who are at increased risk of experiencing syncope during or after the blood sampling [15, 105]. In these patients comfort or distraction may enhance patient response to stress from blood sampling and reduce the risk of syncope.

## IV. Implementation of the guidelines

### Potential barriers and challenges

Successful implementation of the guidelines depends on overcoming any potential barriers or challenges. In order to make a good and feasible implementation plan, one has to first identify all barriers and challenges and carefully consider appropriate solutions (Table 3).

Potential barriers and challenges at the individual level which might compromise successful implementation of this recommendation are the resistance of an individual to change, language barrier, the lack of knowledge, awareness and understanding. Finally, even if there is a positive attitude towards a change, such change could be difficult if there is nobody who is responsible to manage the change or said responsible individual has some other priorities.

Barriers and challenges at the level of the hospital could be of a financial nature. There could also be issues such as the lack of staff who could take over the responsibility to manage the change. Certainly, a change would be difficult if it is deemed as low priority to hospital management.

There are also several possible barriers which could arise at the national level. As is the case at the level of the individual hospital, possible barriers at the national level could be the lack of awareness and understanding about the necessity for implementing the recommendation as well as the lack of a professional entity who could take over the responsibility to manage the change. Also, in some countries there is more than one professional group whose members are involved in the blood sampling process. The existence of such groups might be an obstacle towards the successful implementation of the

**Table 3:** Potential barriers and challenges that need to be overcome for the successful implementation of the guidelines and recommendations.

Barriers and challenges	Solutions
1. Individual <ul style="list-style-type: none"> <li>a. The resistance of an individual to change</li> <li>b. Language barrier</li> <li>c. The lack of knowledge, awareness and understanding about the necessity for implementing the recommendation</li> </ul>	<ul style="list-style-type: none"> <li>a. Change management (shared vision and team work)</li> <li>b. Translate document in local language</li> <li>c. Education</li> </ul>
2. At the level of the hospital <ul style="list-style-type: none"> <li>a. Financial reasons</li> <li>b. The lack of staff who could take over the responsibility to manage the change</li> <li>c. A change is deemed as low priority to hospital management</li> </ul>	<ul style="list-style-type: none"> <li>a. Demonstrate the cost of poor quality to the hospital management</li> <li>b. Identify hospital “ambassador” and build a team</li> <li>c. Present the benefits to the hospital management (savings, patient safety, hospital prestige, etc.)</li> </ul>
3. At the national level <ul style="list-style-type: none"> <li>a. The lack of awareness and understanding about the necessity for implementing the recommendation</li> <li>b. The lack of a professional entity who could take over the responsibility to manage the change</li> <li>c. There is more than one professional group whose members are involved in the blood sampling process</li> <li>d. Recommendations are supported only if they come from a national regulatory body</li> <li>e. The existing national legislation is in conflict with this document</li> <li>f. The recommendation is difficult to implement if it is not officially endorsed or even included in some internationally recognized regulatory document (such as CLSI, ISO, etc.)</li> </ul>	<ul style="list-style-type: none"> <li>a. Identify national “ambassador”</li> <li>b. Establish the national working group for preanalytical phase</li> <li>c. Multidisciplinary collaboration of all stakeholders</li> <li>d. Engage with national regulatory bodies</li> <li>e. Adapt recommendation to local rules and regulations</li> <li>f. EFLM to liaise with international regulatory bodies</li> </ul>

recommendations, if they do not agree to work together. In some countries, recommendations are supported only if they come from a regulatory body. Finally, if the existing national legislation is in conflict with this document, this could pose a considerable difficulty to the implementation of this recommendation.

It could also be that some countries and national associations would find it difficult to implement the recommendation if it is not officially endorsed or even included in some internationally recognized regulatory document (such as CLSI, ISO, etc.).

Given all the mentioned difficulties in finding appropriate communication channels or targeting responsible entities in each country, it can indeed be a great challenge for all EFLM and COLABIOCLI members to accept and implement this recommendation. We therefore propose a framework for a successful implementation of

this recommendation and hope that it might facilitate the implementation process, wherever necessary.

**Framework for a successful implementation of this recommendation**

Necessary requirements for the successful implementation of this recommendation are outlined in the Table 4. In the text below each requirement and its importance are discussed.

There are many ways to deal with the resistance of an individual to a change [106]. We believe that the majority of medical staff are highly concerned with patient safety and well-being. Therefore, their resistance to learn and adopt new blood sampling procedure is basically caused by their lack of understanding of the potential harm to the

**Table 4:** Framework for a successful implementation of EFLM-COLABIOCLI recommendation for venous blood sampling.

Education of the staff	<ul style="list-style-type: none"> <li>– Available already during formal education</li> <li>– Available to all newly employed staff</li> <li>– Available periodically (every 3 years at minimum)</li> <li>– e-learning mode preferable</li> <li>– “Train the trainers” system established</li> <li>– Knowledge test is used prior and after education</li> </ul>
Practical training of the staff	<ul style="list-style-type: none"> <li>– Available already during formal education</li> <li>– Available to all newly employed staff</li> <li>– Available periodically (every 3 years at minimum)</li> <li>– Preferably provided in the laboratory outpatient unit</li> <li>– At least 1 week (at least 100 blood collections) long</li> </ul>
Certification of staff involved in blood sampling	<ul style="list-style-type: none"> <li>– Applies to all who are involved in blood sampling</li> <li>– Granted to new members of staff after successful completion of:                             <ul style="list-style-type: none"> <li>a) Initial education and training</li> <li>b) Knowledge testing and observational audit</li> </ul> </li> <li>– Periodical re-certification</li> </ul>
Auditing of the blood sampling procedure	<ul style="list-style-type: none"> <li>– Periodical auditing system is established</li> <li>– Re-training is done as a corrective measure</li> <li>– Audit is done (observational) using the structured checklist</li> <li>– During audit at least 20 blood collections, performed by at least three different phlebotomists are observed</li> <li>– Quality indicators are used to monitor the sample quality</li> <li>– Quality indicators are used to act upon and initiate corrective measure</li> </ul>
Hospital team responsible for the implementation	<ul style="list-style-type: none"> <li>– There is a hospital “ambassador”</li> <li>– There is a team of key hospital stakeholders</li> </ul>
National societies	<ul style="list-style-type: none"> <li>– There is a national “ambassador”</li> <li>– There is a working group for preanalytical phase in the national society</li> <li>– The recommendation is translated to the local language</li> <li>– Key stakeholders are identified</li> <li>– The implementation is done in collaboration with key stakeholders</li> <li>– Regulatory and governmental bodies support and endorse the implementation activities</li> <li>– All national rules and recommendations take precedence over this document; there is a mechanism to agree on the modifications</li> <li>– Editors of national journals assist by raising awareness</li> </ul>

patient or themselves which may arise as a consequence of non-adherence to the recommended procedure. By educating staff about potential risks to the patient, caused by poor blood sampling procedure, awareness is raised about the necessity to adhere to the recommended procedure [107–109]. Education increases the level of confidence and improves quality of procedures [110]. Nevertheless, the effects are usually short-term and this is why education should be continuously repeated [111].

There is a low level of knowledge and understanding of some basic preanalytical issues among students in biomedicine (medical school, pharmacy, veterinary medicine) [1, 112]. The education about blood sampling procedure should therefore be available to medical staff already during their formal education to become qualified (theoretical and practical). As different professions are involved in blood sampling in different European countries, the professions who would need to receive such education vary from country to country [113].

Education about blood sampling procedure should also be available to all newly employed medical staff involved in blood sampling. Also, besides education, which is mostly theoretical, newly employed staff should undergo a practical training of the blood collection procedure. Practical training should preferably be offered in the laboratory outpatient unit, during the period of 1 week during which a new staff member should perform at least 100 blood collections, under the supervision of the responsible staff. An observational audit should be done during the first five and last five collections, to assess the level of compliance with the recommended procedure and identify potential deviations.

The above stated numbers of blood collections and duration of the practical training are a recommendation for minimum criteria. These criteria are a consensus opinion based on experience and expertise of the authors of this document. We do recognize that the minimum number of blood collections may depend on the institution, the level of skills and experience of the trainee, complexity of intended patient category, etc. It is therefore the responsibility of the educators and trainers that a minimal demonstrable standard of phlebotomy experience and knowledge is achieved.

We recommend that each institution establishes its own system of certification of staff involved in the blood sampling procedure. Certification should be granted to all new members of staff only after successful completion of initial education and training. Knowledge testing and an observational audit are suggested as a requirement for a certificate. To obtain a certificate, a member of the staff should successfully pass the knowledge test. We

recommend 80% of the correct replies, as a success criterion, but it is completely up to the institution to define its minimal standard.

We also recommend that each healthcare institution has a system of continuous auditing, re-training and re-certification for all staff members. We recommend that auditing is done in the form of observational audit using the standardized structured auditing checklist (Table 5). An observational audit should be done periodically in each clinical department at least once per year. During each observational audit, a sufficient number of phlebotomies and phlebotomists should be observed. We recommend that at least 20 blood collections, performed by at least three different phlebotomists (at least three per each phlebotomist) should be observed during each audit. Again, as already stated, it is completely up to the institution to define its minimal standard.

Periodical education (theoretical and practical) should be made available to all staff members after every 3 years at minimum. This education could even be organized as e-learning, if resources are available. As education and training could be time demanding and in settings where human resources are limited, we recommend that a system is established to “train the trainers”, meaning that at each department there is a member of the medical staff (chief nurse of the department) responsible for education, training and auditing of the staff.

We recommend that a knowledge test is used to assess the level of knowledge and understanding as well as to raise awareness of the staff prior to education. Also, we recommend that a knowledge test is used to assess the level of knowledge and awareness of the staff after the education. The knowledge test should assess the understanding about the below listed issues and facts:

- most frequent errors in the preanalytical phase
- the impact of preanalytical errors on the quality of the sample and patient outcome
- how to properly prepare a patient for blood sampling?
- how is fasting defined and why is it important?
- proper patient ID and tube labeling procedure
- tube types, additives
- the order of draw
- the use of tourniquet
- adequate mixing procedure
- why blood-to-additive ratio matters?
- hemolysis – causes and consequences
- clotting – causes and consequences
- patient and healthcare worker safety

Quality indicators are efficient tools for obtaining information about the risk of errors, error frequencies and their

**Table 5:** EFLM-COLABIOCLI venous blood collection observation form.

Observer name:						
Ward/department <sup>a</sup> :						
Date of collection:						
Phlebotomist name/ID:						
Blood collection number	Collection 1		Collection 2		Collection 3	
Question 1. Did the collector properly identify the patient?	yes	no	yes	no	yes	no
Question 2. Did the collector verify that the patient is fasting and properly prepared for phlebotomy?	yes	no	yes	no	yes	no
Question 3. Did the collector obtain all supplies necessary prior to collection?	yes	no	yes	no	yes	no
Question 4. Were the tubes labeled in the presence of the patient?	yes	no	yes	no	yes	no
Question 5. Did the collector put on a new, fresh pair of gloves?	yes	no	yes	no	yes	no
Question 6. Was the tourniquet placed four finger widths (10 cm) above the venipuncture site?	yes	no	yes	no	yes	no
Question 7. Was a suitable venipuncture site selected according to the recommended practice?	yes	no	yes	no	yes	no
Question 8. Was the venipuncture site cleaned properly and not touched after it had been cleaned?	yes	no	yes	no	yes	no
Question 9. Did the collector release the tourniquet when blood flow commenced?	yes	no	yes	no	yes	no
Question 10. Was the first tube (and all subsequent tubes) immediately inverted once gently?	yes	no	yes	no	yes	no
Question 10. Did the collector follow the correct order of draw?	yes	no	yes	no	yes	no
Question 12. Was the safety feature in the blood collection system activated immediately?	yes	no	yes	no	yes	no
Question 13. Was the needle/collection system safely and immediately disposed?	yes	no	yes	no	yes	no
Question 14. Did the collector place a clean gauze over the venipuncture site?	yes	no	yes	no	yes	no
Question 15. Was the patient told to apply pressure until the bleeding has stopped and not to bend the arm?	yes	no	yes	no	yes	no
Question 16. Were all sample tubes mixed for additional 4 times?	yes	no	yes	no	yes	no
Question 17. Did the collector remove his/her gloves once the phlebotomy was completed?	yes	no	yes	no	yes	no
Question 18. Was the patient advised to rest for 5 min to ensure bleeding had stopped before leaving the phlebotomy unit?	yes	no	yes	no	yes	no

<sup>a</sup>Additional generic information related to the institution might be necessary, to properly identify phlebotomist and institutional unit. This will depend on institutional policy and organization as well as on some particular local circumstances. Exclusion criteria: Patients should be conscious, >18 years and blood should not be taken via a catheter. Guide: Use one form per phlebotomist. Each phlebotomist should be monitored during three subsequent phlebotomies.

distribution throughout the total testing process [114]. We recommend that quality indicators are used to monitor the quality of samples received in the laboratory [115–117]. Laboratories are recommended to monitor the frequency of under-filled tubes, clotted samples, sample hemolysis, ID errors, etc. as they are a good tool to detect certain “spikes” and point to some specific problems during the blood collection procedure. The choice of the quality indicators to be used will depend on the local requirements and particular problems and issues at the level of each hospital. Quality indicators should be used to act upon them and correct the issues.

To overcome the language barrier, the recommendation should be translated to the local language and made available to all involved in the blood sampling process. We encourage national societies to assist in the translation of this document.

As regards the ways to overcome barriers at the level of the hospital, one has to be able to present the benefits of the implementation of this recommendation, such as the cost of poor sample quality, potential savings, reduction of patient harm or improvement of patient safety and satisfaction [118, 119]. Furthermore, it has been demonstrated that adherence to the recommended blood

collection procedure minimizes the risk of patient harm and frequency of unsuitable samples [120]. This important safety aspect needs to be demonstrated to the hospital management. Finally, hospital management is likely to be interested in any intervention which could potentially be regarded as a matter of prestige among similar institutions.

For successful implementation of the recommendation, there should be a member of the staff who should be responsible to manage the change at the level of the hospital (a so-called: “ambassador”). This person should have time dedicated for this task.

Also, this person should have a team consisting of several key stakeholders in the hospital, such as the chief nurse and possibly representatives from the:

- laboratory
- clinical staff (medical doctors)
- laboratory technicians
- epidemiologists
- department for hospital infections and worker safety
- quality department
- top hospital management

This team should meet on a regular basis and discuss and plan strategy for successful implementation and continuous improvement.

At the national level, there should also be an “ambassador” who will take the lead in the process of implementation of this recommendation. To facilitate the implementation there should be a working group for the preanalytical phase or some other entity which will be responsible for educational interventions and raising the awareness among all stakeholders and professions (of the same or different background and level of education) involved in blood sampling about the necessity for the implementation of the recommendation. National journals and their editors are also encouraged to raise awareness about preanalytical phase and venous blood sampling in particular, by offering their journal as an efficient and powerful vehicle for sharing knowledge and information [121–123]. The implementation process should be done as a joint effort in close multidisciplinary collaboration of all stakeholders at the national level. National “ambassadors” are responsible to identify and recruit key stakeholders such as national nursing associations, professional societies in laboratory medicine and preferably even patients.

It is highly advisable to involve regulatory bodies, such as professional chambers, associations, national regulatory entities and even governmental bodies like Ministry of Health to support and endorse the implementation activities.

If some national rules are in conflict with this document, there should be a mechanism to agree on the modification of this recommendation at the national level and accept the revised version for implementation.

## Conclusions

The EFLM WG-PRE as the leading professional entity involved in preanalytical phase feels responsible to provide a framework for a successful implementation of this document at the European level [124, 125]. It is our aim to encourage the European Association for Accreditation to endorse this document as a standard and encourage its use at the national level in each European country during accreditation assessments.

To facilitate the implementation EFLM WG-PRE has prepared following tools:

1. a power point presentation describing some basic issues related to venous blood sampling and the entire procedure (to be used during the education of staff)
2. video describing the entire procedure (to be used during the education of staff)
3. a knowledge test to assess the level of knowledge and raise awareness of the staff prior and after the education
4. a checklist to be used for auditing the blood sampling procedure during periodical observational audits (Table 5)
5. posters with a cartoon describing the entire procedure (to be used at blood collection facilities)

These tools are freely available at the EFLM website ([www.eflm.eu](http://www.eflm.eu)) under EFLM Committees/Science/WG:Preanalytical Phase, under Resources/Educational Material. Professionals are encouraged to download and use these tools to implement the recommended procedure for venous blood collection and establish a quality system in place to maintain and continuously improve the quality of the procedure.

**Author contributions:** All the authors have accepted responsibility for the entire content of this submitted manuscript and approved submission.

**Research funding:** None declared.

**Employment or leadership:** None declared.

**Honorarium:** None declared.

**Competing interests:** The funding organization(s) played no role in the study design; in the collection, analysis, and interpretation of data; in the writing of the report; or in the decision to submit the report for publication.

## References

1. Simundic AM, Cornes M, Grankvist K, Lippi G, Nybo M, Kovalevskaia S, et al. Survey of national guidelines, education and training on venous blood collection in 28 European countries: an original report by the European Federation of Clinical Chemistry and Laboratory Medicine (EFLM) working group for the preanalytical phase (WG-PA). *Clin Chem Lab Med* 2013;51:1585–93.
2. Lippi G, Cervellin G, Mattiuzzi C. Critical review and meta-analysis of spurious hemolysis in blood samples collected from intravenous catheters. *Biochem Med (Zagreb)* 2013;23:193–200.
3. Mrazek C, Simundic AM, Wiedemann H, Kraher F, Felder TK, Kipman U, et al. The relationship between vacuum and hemolysis during catheter blood collection: a retrospective analysis of six large cohorts. *Clin Chem Lab Med* 2017;55:1129–34.
4. Heiligers-Duckers C, Peters NA, van Dijk JJ, Hoeijmakers JM, Janssen M. Low vacuum and discard tubes reduce hemolysis in samples drawn from intravenous catheters. *Clin Biochem* 2013;46:1142–4.
5. ISO/TS 15189:2012 Medical laboratories – Requirements for quality and competence.
6. ISO/TS 20658:2017 Medical laboratories – Requirements for collection, transport, receipt, and handling of samples.
7. Simundic AM, Church S, Cornes MP, Grankvist K, Lippi G, Nybo M, et al. Compliance of blood sampling procedures with the CLSI H3-A6 guidelines: an observational study by the European Federation of Clinical Chemistry and Laboratory Medicine (EFLM) working group for the preanalytical phase (WG-PRE). *Clin Chem Lab Med* 2015;53:1321–31.
8. Clinical Laboratory Standards Institute. GP41: procedures for collection of diagnostic blood specimens by venipuncture; approved guideline, 7th ed. CLSI document GP41. Wayne, PA: Clinical and Laboratory Standards Institute, 2007.
9. World Health Organization. WHO guidelines on drawing blood. [http://whqlibdoc.who.int/publications/2010/9789241599221\\_eng.pdf](http://whqlibdoc.who.int/publications/2010/9789241599221_eng.pdf). Accessed: 11 Jan 2013.
10. Guyatt GH, Oxman AD, Kunz R, Falck-Ytter Y, Vist GE, Liberati A, et al. Going from evidence to recommendations. *Br Med J* 2008;336:1049–51.
11. <http://www.uptodate.com/home/grading-guide#gradingrecomendations>. Accessed: June 2018.
12. EFLM Procedure Manual v1.15, April 2017; Accessed: 9 Jun 2018, under Official Documents/Rules and regulations at: <https://www.eflm.eu/site/page/a/1056>.
13. American College of Obstetricians and Gynecologists Committee on Health Care for Underserved Women, Committee on Patient Safety, Quality Improvement. ACOG Committee Opinion No. 587: effective patient-physician communication. *Obstet Gynecol* 2014;123:389–93.
14. Ha JF, Longnecker N. Doctor-patient communication: a review. *Ochsner J* 2010;10:38–43.
15. France CR, France JL, Himawan LK, Stephens KY, Frame-Brown TA, Venable GA, et al. How afraid are you of having blood drawn from your arm? A simple fear question predicts vasovagal reactions without causing them among high school donors. *Transfusion* 2013;53:315–21.
16. Simundic AM, Nikolac N, Guder W. Preanalytical variation and preexamination processes. In: Rifai N, Horvath R, Wittwer C, editors. *Tietz textbook of clinical chemistry and molecular diagnostics*, 6th ed. St. Louis, Missouri, USA: Elsevier, 2018:81–120.
17. Lippi G, Salvagno GL, Lima-Oliveira G, Danese E, Favaloro EJ, Guidi GC. Influence of posture on routine hemostasis testing. *Blood Coagul Fibrinolysis* 2015;26:716–9.
18. Lippi G, Salvagno GL, Lima-Oliveira G, Brocco G, Danese E, Guidi GC. Postural change during venous blood collection is a major source of bias in clinical chemistry testing. *Clin Chim Acta* 2015;440:164–8.
19. Lippi G, Cervellin G. Acutely developing, spurious anemia without actual blood loss. A paradigmatic case report. *Biochem Med* 2017;27:421–5.
20. Lima-Oliveira G, Guidi GC, Salvagno GL, Danese E, Montagnana M, Lippi G. Patient posture for blood collection by venipuncture: recall for standardization after 28 years. *Rev Bras Hematol Hemoter* 2017;39:127–32.
21. van Dongen-Lases E, Cornes MP, Grankvist K, Ibarz M, Kristensen GB, Lippi G, et al. Patient identification and tube labelling – a call for harmonisation on behalf of the Working Group for Preanalytical Phase (WG-PRE), European Federation of Clinical Chemistry and Laboratory Medicine (EFLM). *Clin Chem Lab Med* 2016;54:1141–5.
22. Simundic AM, Cornes M, Grankvist K, Lippi G, Nybo M. Standardization of collection requirements for fasting samples. For the Working Group on Preanalytical Phase (WG-PA) of the European Federation of Clinical Chemistry and Laboratory Medicine (EFLM). *Clin Chim Acta* 2014;432:33–7.
23. Lima-Oliveira G, Volanski W, Lippi G, Picheth G, Guidi GC. Preanalytical phase management: a review of the procedures from patient preparation to laboratory analysis. *Scand J Clin Lab Invest* 2017;77:153–63.
24. Simundic AM, Dorotić A, Fumic K, Gudasic-Vrdoljak J, Kackov S, Klenkar K, et al. Patient preparation for laboratory testing: recommendation of the Croatian Society of Medical Biochemistry and Laboratory Medicine. *Biochem Med* 2018. In press.
25. Nikolac N, Supak-Smolcic V, Simundic AM, Celap I. Croatian Society of Medical Biochemistry and Laboratory Medicine: national recommendations for venous blood sampling. *Biochem Med* 2013;23:242–54.
26. Montagnana M, Danese E, Salvagno GL, Lippi G. Short-term effect of dark chocolate consumption on routine haemostasis testing. *Int J Food Sci Nutr* 2017;68:613–6.
27. Lippi G, Lima-Oliveira G, Salvagno GL, Montagnana M, Gelati M, Picheth G, et al. Influence of a light meal on routine haematological tests. *Blood Transfus* 2010;8:94–9.
28. Lima-Oliveira G, Salvagno GL, Lippi G, Gelati M, Montagnana M, Danese E, et al. Influence of a regular, standardized meal on clinical chemistry analytes. *Ann Lab Med* 2012;32:250–6.
29. Lima-Oliveira G, Salvagno GL, Lippi G, Danese E, Gelati M, Montagnana M, et al. Could light meal jeopardize laboratory coagulation tests? *Biochem Med (Zagreb)* 2014;24:343–9.
30. Simundic AM, Filipi P, Vrtaric A, Miler M, Nikolac Gabaj N, Kocsis A, et al. Patient's knowledge and awareness about the effect of the over-the-counter (OTC) drugs and dietary supplements on laboratory test results: a survey in 18 European countries. *Clin Chem Lab Med*. 2018, in press.
31. Perovic A, Nikolac N, Braticovic NM, Milcic A, Sobocanec S, Balog T, et al. Does recreational scuba diving have clinically significant effect on routine haematological parameters? *Biochem Med* 2017;27:325–31.
32. Danese E, Salvagno GL, Tarperi C, Negrini D, Montagnana M, Festa L, et al. Middle-distance running acutely influences the

- concentration and composition of serum bile acids. Potential implications for cancer risk? *Oncotarget* 2017;8:52775–82.
33. Corsetti R, Lombardi G, Barassi A, Lanteri P, Colombini A, D'Eril GM, et al. Cardiac indexes, cardiac damage biomarkers and energy expenditure in professional cyclists during the Giro d'Italia 3-weeks stage race. *Biochem Med* 2012;22:237–46.
  34. Rasaiah B, Hoag G. Guidelines for a venous blood collection chair. *Can Med Assoc J* 1992;146:108–9.
  35. Lippi G, Cornes MP, Grankvist K, Nybo M, Simundic AM. European Federation of Clinical Chemistry and Laboratory Medicine (EFLM) Working Group for Preanalytical Phase (WG-PRE) opinion paper: local validation of blood collection tubes in clinical laboratories. *Clin Chem Lab Med* 2016;54:755–60.
  36. Bostic G, Thompson R, Atanasoski S, Canlas C, Ye H, Kolins M, et al. Quality improvement in the coagulation laboratory: reducing the number of insufficient blood draw specimens for coagulation testing. *Lab Med* 2015;46:347–55.
  37. Domingos MC, Médaille C, Concordet D, Briend-Marchal A. Is it possible to use expired tubes for routine biochemical analysis in dogs? *Vet Clin Pathol* 2012;41:266–71.
  38. Verbeek JH, Ijaz S, Mischke C, Ruotsalainen JH, Mäkelä E, Neuvonen K, et al. Personal protective equipment for preventing highly infectious diseases due to exposure to contaminated body fluids in healthcare staff. *Cochrane Database Syst Rev* 2016;4:CD011621.
  39. Kinlin LM, Mittleman MA, Harris AD, Rubin MA, Fisman DN. Use of gloves and reduction of risk of injury caused by needles or sharp medical devices in healthcare workers: results from a case-crossover study. *Infect Control Hosp Epidemiol* 2010;31:908–17.
  40. Mast ST, Woolwine JD, Gerberding JL. Efficacy of gloves in reducing blood volumes transferred during simulated needlestick injury. *J Infect Dis* 1993;168:1589–92.
  41. De Carli G, Abiteboul D, Puro V. The importance of implementing safe sharps practices in the laboratory setting in Europe. *Biochem Med (Zagreb)* 2014;24:45–56.
  42. Bhargava A, Mishra B, Thakur A, Dogra V, Loomba P, Gupta S. Assessment of knowledge attitude and practices among healthcare workers in a tertiary care hospital on needle stick among injury. *Int J Health Care Qual Assur* 2013;26:549–58.
  43. Self WH, Mickanin J, Grijalva CG, Grant FH, Henderson MC, Corley G, et al. Reducing blood culture contamination in community hospital emergency departments: a multicenter evaluation of a quality improvement intervention. *Acad Emerg Med* 2014;21:274–82.
  44. Self WH, Speroff T, Grijalva CG, McNaughton CD, Ashburn J, Liu D, et al. Reducing blood culture contamination in the emergency department: an interrupted time series quality improvement study. *Acad Emerg Med* 2013;20:89–97.
  45. Mansouri M, Tidley M, Sanati KA, Roberts C. Comparison of blood transmission through latex and nitrile glove materials. *Occup Med* 2010;60:205–10.
  46. Wittman A, Kralj N, Köver J, Gasthaus K, Lerch H, Hofmann F. Comparison of 4 different types of surgical gloves used for preventing blood contact. *Infect Control Hosp Epidemiol* 2010;31:498–502.
  47. Pittet D, Allegranzi B, Sax H, Dharan S, Pessoa-Silva CL, Donaldson L, et al. Evidence-based model for hand transmission during patient care and the role of improved practices. *Lancet Infect Dis* 2006;6:641–52.
  48. Dukic K, Zoric M, Pozaic P, Starcic J, Culjak M, Saracevic A, et al. How compliant are technicians with universal safety measures in medical laboratories in Croatia? – a pilot study. *Biochem Med* 2015;25:386–92.
  49. Lima-Oliveira G, Lippi G, Salvagno GL, Montagnana M, Picheth G, Guidi GC. Impact of the venous blood collection training based on CLSI/NCCLS H03–A6 – procedures for the collection of diagnostic blood specimens by venipuncture. *Biochem Med (Zagreb)* 2012;22:342–51.
  50. Culjak M, Gveric Grginic A, Simundic AM. Bacterial contamination of reusable venipuncture tourniquets in tertiary-care hospital. *Clin Chem Lab Med* 2018; doi: 10.1515/cclm-2017-0994.
  51. Mehmood Z, Muhammad Mubeen S, Shehzad Afzal M, Husain Z. Potential risk of cross-infection by tourniquets: a need for effective control practices in Pakistan. *Int J Prev Med* 2014;5:1119–24.
  52. Pinto AN, Phan T, Sala G, Cheong EY, Siarakas S, Gottlieb T. Reusable venesection tourniquets: a potential source of hospital transmission of multiresistant organisms. *Med J Aust* 2011;195:276–9.
  53. Nikolac N, Lenicek Krleza J, Simundic AM. Preanalytical external quality assessment of the Croatian Society of Medical Biochemistry and Laboratory Medicine and CROQALM: finding undetected weak spots. *Biochem Med* 2017;27:131–43.
  54. Lima-Oliveira G, Lippi G, Salvagno GL, Montagnana M, Manguera CL, Sumita NM, et al. New ways to deal with known preanalytical issues: use of transilluminator instead of tourniquet for easing vein access and eliminating stasis on clinical biochemistry. *Biochem Med* 2011;21:152–9.
  55. Lima-Oliveira G, Lippi G, Salvagno GL, Montagnana M, Scartezini M, Guidi GC, et al. Transillumination: a new tool to eliminate the impact of venous stasis during the procedure for the collection of diagnostic blood specimens for routine haematological testing. *Int J Lab Hematol* 2011;33:457–62.
  56. Lima-Oliveira G, Salvagno GL, Lippi G, Montagnana M, Scartezini M, Picheth G, et al. Elimination of the venous stasis error for routine coagulation testing by transillumination. *Clin Chim Acta* 2011;412:1482–4.
  57. Don BR, Sebastian A, Cheitlin M, Christiansen M, Schambelan M. Pseudohyperkalemia caused by fist clenching during venous blood collection. *N Engl J Med* 1990;322:1290–2.
  58. Seimiya M, Yoshida T, Sawabe Y, Sogawa K, Umemura H, Matsushita K, et al. Reducing the incidence of pseudohyperkalemia by avoiding making a fist during venous blood collection: a quality improvement report. *Am J Kidney Dis* 2010;56:686–92.
  59. Ialongo C, Bernardini S. Phlebotomy, a bridge between laboratory and patient. *Biochem Med* 2016;26:17–33.
  60. Loh TP, Sethi SK. A multidisciplinary approach to reducing spurious hyperkalemia in hospital outpatient clinics. *J Clin Nurs* 2015;24:2900–6.
  61. Lima-Oliveira G, Guidi GC, Salvagno GL, Lippi G. The impact of fist clenching and its maintenance during venipuncture on routine hematology testing. *J Clin Lab Anal* 2017;31. doi: 10.1002/jcla.22108.
  62. Lima-Oliveira G, Guidi GC, Salvagno GL, Brocco G, Danese E, Lippi G. Estimation of the imprecision on clinical chemistry testing due to fist clenching and maintenance during venipuncture. *Clin Biochem* 2016;49:1364–7.
  63. Putz R, Pabst R, editors. Sobotta: atlas of human anatomy, 20th ed. Munich, DE: Urban & Schwarzenberg/Elsevier, 1993.

64. Horowitz SH. Venipuncture-induced causalgia: anatomic relations of upper extremity superficial veins and nerves, and clinical considerations. *Transfusion* 2000;40:1036–40.
65. Ramos JA. Venipuncture-related lateral antebrachial cutaneous nerve injury: what to know? *Braz J Anesthesiol* 2014;64:131–3.
66. Seifert H, Abele-Horn M, Fätkenheuer G, Shah PM. Mikrobiologische-infektiologische Qualitätsstandards (MiQ) – Blutkulturdiagnostik, Urban&Fischer 2007, S.16–27 (in German).
67. Anforderungen an die Hygiene bei Punktionen und Injektionen Empfehlung der Kommission für Krankenhaushygiene und Infektionsprävention beim Robert Koch-Institut (RKI). *Bundesgesundheitsbl* 2011;54:1135–44. (in German).
68. Patel TG, Shukla RV, Gupte SC. Impact of donor arm cleaning with different aseptic solutions for prevention of contamination in blood bags. *Indian J Hematol Blood Transfus* 2013;29:17–20.
69. Ibáñez-Cervantes G, Bello-López JM, Fernández-Sánchez V, Domínguez-Mendoza CA, Acevedo-Alfaro LI. Prevalence of bacterial contamination in platelet concentrates at the National Center of Blood Transfusion (Mexico). *Transfus Clin Biol* 2017;24:56–61.
70. Pendlington RU, Whittle E, Robinson JA, Howes D. Fate of ethanol topically applied to skin. *Food Chem Toxicol* 2001;39:169–74.
71. Salvagno GL, Danese E, Lima-Oliveira G, Guidi GC, Lippi G. Avoidance to wipe alcohol before venipuncture is not a source of spurious hemolysis. *Biochem Med* 2013;23:201–5.
72. Lippi G, Simundic AM, Musile G, Danese E, Salvagno G, Tagliaro F. The alcohol used for cleansing the venipuncture site does not jeopardize blood and plasma alcohol measurement with head-space gas chromatography and an enzymatic assay. *Biochem Med* 2017;27:398–403.
73. Hadaway LC, Millam DA. On the road to successful I.V. starts. *Nursing* 2005;35(Suppl On):1–14; quiz 14–6.
74. Cornes M, van Dongen-Lases E, Grankvist K, Ibarz M, Kristensen G, Lippi G, et al. Order of blood draw: opinion paper by the European Federation for Clinical Chemistry and Laboratory Medicine (EFLM) Working Group for the Preanalytical Phase (WG-PRE). *Clin Chem Lab Med* 2017;55:27–31.
75. Smock KJ, Crist RA, Hansen SJ, Rodgers GM, Lehman CM. Discard tubes are not necessary when drawing samples for specialized coagulation testing. *Blood Coagul Fibrinolysis* 2010;21:279–82.
76. Lippi G, Guidi GC. Effect of specimen collection on routine coagulation assays and D-dimer measurement. *Clin Chem* 2004;50:2150–2.
77. Sulaiman RA, Cornes MP, Whitehead S, Othonos N, Ford C, Gama R. Effect of order of draw of blood samples during venous blood collection on routine biochemistry results. *J Clin Pathol* 2011;64:1019–20.
78. Salvagno G, Lima-Oliveira G, Brocco G, Danese E, Guidi GC, Lippi G. The order of draw: myth or science? *Clin Chem Lab Med* 2013;51:2281–5.
79. Cornes MP, Ford C, Gama R. Spurious hyperkalaemia due to EDTA contamination: common and not always easy to identify. *Ann Clin Biochem* 2008;45:601–3.
80. Lima-Oliveira G, Lippi G, Salvagno GL, Montagnana M, Picheth G, Guidi GC. Incorrect order of draw could be mitigate the patient safety: a phlebotomy management case report. *Biochem Med (Zagreb)* 2013;23:218–23.
81. Sharratt CL, Gilbert CJ, Cornes MP, Ford C, Gama R. EDTA sample contamination is common and often undetected, putting patients at unnecessary risk of harm. *Int J Clin Pract* 2009;63:1259–62.
82. Cadamuro J, Felder TK, Oberkofler H, Mrazek C, Wiedemann H, Haschke-Becher E. Relevance of EDTA carryover during blood collection. *Clin Chem Lab Med* 2015;53:1271–8.
83. Berg JE, Ahee P, Berg JD. Variation in venous blood collection techniques in emergency medicine and the incidence of haemolysed samples. *Ann Clin Biochem* 2011;48(Pt 6):562–5.
84. Lippi G, Salvagno GL, Montagnana M, Brocco G, Guidi GC. Influence of short-term venous stasis on clinical chemistry testing. *Clin Chem Lab Med* 2005;43:869–75.
85. Lippi G, Salvagno GL, Montagnana M, Guidi GC. Short-term venous stasis influences routine coagulation testing. *Blood Coagul Fibrinolysis* 2005;16:453–8.
86. Lippi G, Salvagno GL, Montagnana M, Franchini M, Guidi GC. Venous stasis and routine hematologic testing. *Clin Lab Haematol* 2006;28:332–7.
87. Lima-Oliveira G, Lippi G, Salvagno GL, Montagnana M, Gelati M, Volanski W, et al. Effects of vigorous mixing of blood vacuum tubes on laboratory test results. *Clin Biochem* 2013;46:250–4.
88. Karlsson J, Helmersson-Karlqvist J, Larsson A. Delayed mixing of vacuum tubes clearly affects platelet counts but not haemoglobin concentration and prothrombin time (INR) results. *Int J Lab Hematol* 2013;35:15–7.
89. Clinical Laboratory Standards Institute. Collection, Transport, and Processing of Blood Specimens for Testing Plasma-Based Coagulation Assays and Molecular Hemostasis Assays. CLSI H21-A5 document. 5th ed. Wayne, PA: Clinical Laboratory Standards Institute, 2008.
90. Lima-Oliveira G, Lippi G, Salvagno GL, Brocco G, Gaino S, Dima F, et al. Processing of diagnostic blood specimens: is it really necessary to mix primary blood tubes after collection with evacuated tube system? *Biopreserv Biobank* 2014;12:53–9.
91. Parenmark A, Landberg E. To mix or not to mix venous blood samples collected in vacuum tubes? *Clin Chem Lab Med* 2011;49:2061–3.
92. Lippi G, Salvagno GL, Montagnana M, Banfi G, Guidi GC. Evaluation of different mixing procedures for K2 EDTA primary samples on hematological testing. *Lab Med* 2007;38:723–5.
93. Lippi G, Salvagno GL, Montagnana M, Guidi GC. Influence of primary sample mixing on routine coagulation testing. *Blood Coagul Fibrinolysis* 2007;18:709–11.
94. Lippi G, Plebani M. Primary blood tubes mixing: time for updated recommendations. *Clin Chem Lab Med* 2012;50:599–600.
95. Lima-Oliveira G, Lippi G, Salvagno GL, Picheth G, Guidi GC. Laboratory diagnostics and quality of blood collection. *J Med Biochem* 2015;34:288–94.
96. Directive 2010/32/EU – prevention from sharp injuries in the hospital and healthcare sector. <https://osha.europa.eu/es/legislation/directives/council-directive-2010-32-eu-prevention-from-sharp-injuries-in-the-hospital-and-healthcare-sector>. Accessed: 20 Jul 2017.
97. Hansen HC, Harboe H, Drenck NE. Bruising after venepuncture. *Ugeskr Laeger* 1989;151:626–7.
98. Blackmore M. Minimising bruising in the antecubital fossa after venipuncture. *Br Med J (Clin Res Ed)* 1987;295:332.

99. Dyson A, Bogod D. Minimising bruising in the antecubital fossa after venipuncture. *Br Med J (Clin Res Ed)* 1987;294:1659.
100. Godwin PG, Cuthbert AC, Choyce A. Reducing bruising after venepuncture. *Qual Health Care* 1992;1:245–6.
101. Backman C, Zoutman DE, Marck PB. An integrative review of the current evidence on the relationship between hand hygiene interventions and the incidence of health care-associated infections. *Am J Infect Control* 2008;36:333–48.
102. Vissers D, Matthyssen B, Truijien S, Blommaert S, Van De Velde K, Van Gaal L. Fainting and hemolysis during blood sampling in youngsters: prevalence study. *Int J Nurs Stud* 2008;45:760–4.
103. Martens RJ, Geijselaers SL, Stehouwer CD, Henry RM; Maastricht Study Group. Timing of syncope during blood sampling – the Maastricht Study. *Eur J Intern Med* 2017;43:e46–7.
104. Graham DT. Prediction of fainting in blood donors. *Circulation* 1961;23:901–6.
105. France CR, France JL, Kowalsky JM, Ellis GD, Copley DM, Geneser A, et al. Assessment of donor fear enhances prediction of presyncopal symptoms among volunteer blood donors. *Transfusion* 2012;52:375–80.
106. Kotter JP. *Leading change*. Harvard Business Review Press, 1996.
107. Makhumula-Nkhoma N, Whittaker V, McSherry R. Level of confidence in venepuncture and knowledge in determining causes of blood sample haemolysis among clinical staff and phlebotomists. *J Clin Nurs* 2015;24:370–85.
108. Dorotić A, Antončić D, Biljak VR, Nedić D, Beletić A. Hemolysis from a nurses' standpoint—survey from four Croatian hospitals. *Biochem Med (Zagreb)* 2015;25:393–400.
109. Milutinović D, Andrijević I, Ličina M, Andrijević L. Confidence level in venipuncture and knowledge on causes of in vitro hemolysis among healthcare professionals. *Biochem Med (Zagreb)* 2015;25:401–9.
110. Lima-Oliveira G, Lippi G, Salvagno GL, Montagnana M, Picheth G, Guidi GC. Impact of the phlebotomy training based on CLSI/NCCLS H03-A6- procedures for the collection of diagnostic blood specimens by venipuncture. *Biochem Med* 2012;22:342–51.
111. Bölenius K, Lindkvist M, Brulin C, Grankvist K, Nilsson K, Söderberg J. Impact of a large-scale educational intervention program on venous blood specimen collection practices. *BMC Health Serv Res* 2013;13:463.
112. Dukic L, Jokic A, Kules J, Pasalic D. The knowledge and understanding of preanalytical phase among biomedicine students at the University of Zagreb. *Biochem Med* 2016;26:90–7.
113. Simundic AM. Who is doing Phlebotomy in Europe? In: Guder WG, Narayanan S, editors. *Pre-examination procedures in laboratory diagnostics. Preanalytical Aspects and their Impact on the Quality of Medical Laboratory Results*. Berlin, Boston: De Gruyter, 2015.
114. Sciacovelli L, Panteghini M, Lippi G, Sumarac Z, Cadamuro J, Galoro CA, et al. Defining a roadmap for harmonizing quality indicators in Laboratory Medicine: a consensus statement on behalf of the IFCC Working Group “Laboratory Error and Patient Safety” and EFLM Task and Finish Group “Performance specifications for the extra-analytical phases”. *Clin Chem Lab Med* 2017;55:1478–88.
115. Plebani M, Sciacovelli L, Aita A, Chiozza ML. Harmonization of pre-analytical quality indicators. *Biochem Med (Zagreb)* 2014;24:105–13.
116. Plebani M, Sciacovelli L, Aita A, Pelloso M, Chiozza ML. Performance criteria and quality indicators for the pre-analytical phase. *Clin Chem Lab Med* 2015;53:943–8.
117. Plebani M; EFLM Task Force on Performance Specifications for the extra-analytical phases. Performance specifications for the extra-analytical phases of laboratory testing: why and how. *Clin Biochem* 2017;50:550–4.
118. Karcher DS, Lehman CM. Clinical consequences of specimen rejection: a College of American Pathologists Q-Probes analysis of 78 clinical laboratories. *Arch Pathol Lab Med* 2014;138:1003–8.
119. Lippi G, Bonelli P, Cervellin G. Prevalence and cost of hemolyzed samples in a large urban emergency department. *Int J Lab Hematol* 2014;36:e24–6.
120. Ong ME, Chan YH, Lim CS. Reducing blood sample hemolysis at a tertiary hospital emergency department. *Am J Med* 2009;122:1054.e1–6.
121. Simundic AM, Cadamuro J, Cornes J. Biochemia Medica introduces new section: pre-analytical mysteries. *Biochem Med* 2017;27:418–20.
122. Cornes M. Case report of unexpected hypocalcaemia in a slightly haemolysed sample. *Biochem Med (Zagreb)* 2017;27:426–9.
123. Cadamuro J, Wiedemann H, Felder TK, Mrazek C, Kipman U, Hannes O, et al. What/s floating on my plasma? *Biochem Med (Zagreb)* 2017;27:430–3.
124. Lippi G, Simundic AM; European Federation for Clinical Chemistry and Laboratory Medicine (EFLM) Working Group for Preanalytical Phase (WG-PRE). The EFLM strategy for harmonization of the preanalytical phase. *Clin Chem Lab Med* 2017. doi: 10.1515/cclm-2017-0277
125. Cornes MP, Church S, van Dongen-Lases E, Grankvist K, Guimarães JT, Ibarz M, et al. The role of European Federation of Clinical Chemistry and Laboratory Medicine Working Group for Preanalytical Phase in standardization and harmonization of the preanalytical phase in Europe. *Ann Clin Biochem* 2016;53(Pt 5):539–47.

**Supplementary Material:** The online version of this article offers supplementary material (<https://doi.org/10.1515/cclm-2018-0602>).